

STUDY DESIGN ARTICLE

Design of a prospective follow-up study on early parenthood and smoking behaviour during pregnancy in Finnish primary healthcare

MIKAEL O. EKBLAD¹ , HANNA P. WALLIN^{1,2}, MARJUKKA PAJULO³
& PÄIVI E. KORHONEN¹

¹Department of General Practice, University of Turku and Turku University Hospital, Finland, ²Central Satakunta Federation of Municipalities, Finland, and ³Department of Child Psychiatry, University of Turku, Finland

Abstract

Aims: The primary aim of the study is to explore different factors affecting parents' smoking behaviour, and especially how smoking may be connected with individual differences in the psychological process of becoming a parent. In the current paper, we present the study design together with basic information on the study population. **Methods:** The Central Satakunta Maternity and Child Health Clinic (KESALATU) Study is an ongoing prospective follow-up study in primary healthcare of the Satakunta region of southwest Finland. Families were recruited during their first maternity clinic visit between 1 September 2016 and 31 December 2019, and participation will continue until the child is 1.5 years of age. The study combines different sources and types of data: e.g. routine data obtained from primary healthcare clinic records, specific parental self-report data and data from a new exhaled carbon monoxide meter indicating maternal smoking. The data are collected using frequently repeated assessments both during pregnancy and postnatally. The methods cover the following areas of interest: family background factors (including smoking and alcohol use), self-reported parental–foetal/infant attachment and mentalization, self-reported stress, depression and quality of life. **Results:** 589 pregnant women and their partners were asked to participate in the study during the collection time period. The final study population consisted of 248 (42.1%) pregnant women and 160 (27.1%) partners. **Conclusions:** **The new methods and study design have the potential to increase our understanding about the link between early parenting psychology, prenatal psychosocial risk factors and parental health behaviour.**

Keywords: Primary care, maternity clinic, prenatal exposure, smoking, e-cigarettes, pregnancy, attachment, mentalization, depression, anxiety

Introduction

Maternal smoking during pregnancy is the most common harmful exposure during pregnancy compromising foetal and child health. Smoking during pregnancy has been linked with several well-known adverse effects on pregnancy and foetal outcomes including preterm birth, low birth weight and even neonatal morbidity [1,2]. Maternal smoking disturbs significantly the development of many important organs, especially the central nervous system and lungs of the foetus leading to potential long-term adverse effects, which may result in later psychological and psychological problems [2–4].

Maternal second-hand smoke exposure is also known to have adverse effects on child health [5].

Still one in nine of pregnant women continue to smoke in Finland [6]. It is well established that smoking during pregnancy is more common in women who are young, have low educational level, whose partner is a smoker and who have psychiatric problems [4,7]. Economic stress and socioeconomic status across the mother's life course are shown to predict maternal smoking [8]. A study by Härkönen et al. showed that socio-economic disparities in smoking during pregnancy are primarily explained by differences in the mother's level of education [9].

Correspondence: Mikael O. Ekblad, MD, PhD, Department of General Practice, Institute of Clinical Medicine, University of Turku and Turku University Hospital, 20014 Turku, Finland. E-mail: moekbl@utu.fi

Date received 2 December 2020; reviewed 7 May 2021; accepted 13 May 2021

© Author(s) 2021

Article reuse guidelines: sagepub.com/journals-permissions

DOI: 10.1177/14034948211022433

journals.sagepub.com/home/sjp



The women whose partners smoke continue to smoke during pregnancy four times more often compared to women whose partners do not smoke [10].

Finland is one of the first countries to set an ambitious goal to end the use of tobacco and other nicotine products, meaning less than 5% of adults consuming tobacco or nicotine products, by the year 2030 [11]. The prevalence of smoking among adults has been decreasing during the last decades and currently 15% of men and 13% of women smoke daily in Finland [12]. Despite the general positive changes in the prevalence of smoking, it is worthy of notice that a similar decrease has not yet occurred among Finnish pregnant women, and especially not during the early pregnancy phase [6–7].

Parental–foetal/infant attachment and parent mentalization might offer new intervention routes for smoking cessation among the parents-to-be. Prenatal attachment refers to the emotional attachment of a parent to the foetus, which is manifested both in parental behaviour and in the content and amount of mental focusing on the foetus [13]. The relationship between the parent and child starts to develop already when a desire for a child exists [14]. The individual process of becoming a parent for the first time is one of the most intense transitional stages in human life. This requires mental preparation and adjustment, and often brings up ambivalent and vulnerable thoughts [14]. The mental representations, the internalized memories of parental practices and experiences from one's own childhood, become stronger when the mother can feel the movement of the foetus and the baby becomes more real; this starts to strongly shape the quality of the parent–child relationship, that is parent–foetal attachment [15–17].

Mentalization refers to the parent's capability to think of the child as a separate, individual being from early on [18]. A parent who is capable of mentalizing the baby/child is able to accept the necessary changes in daily lifestyle and rhythm that pregnancy and parenthood bring, as well as being able to imagine both the challenging and the most rewarding postnatal situations with the child [19–20]. A strong mentalization capacity contributes to a stronger parent–foetus/infant attachment and strengthens early interaction with the child [21].

The Finnish maternity and child health clinics offer a good opportunity for supporting smoking cessation and the development of parental attachment. The clinics are run by the municipalities in Finland, and the Ministry of Social Affairs and Health regulate the scheme [22]. The maternity and child health clinic visits are free-of-charge for all residents in Finland, thus, in practice, all pregnant women attend the maternity clinics. On average, expectant women

have 11 nurse and/or physician appointments during pregnancy.

There is an urgent need to try to discover how to better enhance smoking cessation among both pregnant women and their partners. Those women who continue to smoke during pregnancy most likely suffer from stronger nicotine dependency and have also other challenges in their life management. This study has the potential to find new intervention routes and to gain knowledge about for whom to provide more comprehensive family support. Conducting research in maternity and child health clinics in a primary healthcare setting is an important way to improve current practices that will lead to better health care.

Overview and aims of the current study

The Central Satakunta Maternity and Child Health Clinic (KESALATU) Study is an ongoing prospective follow-up study on the primary healthcare of the Central Satakunta region of southwest Finland. It was designed to gather comprehensive data on pregnant women and their partners, as well as their children up to the age of 1.5 years. The data are collected using different sources of information, for example maternity and child healthcare clinic records, parental self-reports and a new exhaled breath carbon monoxide (CO) meter assessing amount of tobacco smoking from exhaled air. The primary aims of the study are presented in Table I.

Material and methods

Study region

The Central Satakunta Health Federation of Municipalities is responsible for organizing the maternity and child health clinics in the study area. It includes four municipalities, participating in the current study (Harjavalta, Nakkila, Kokemäki and Eurajoki), with a total population of around 30,000 inhabitants.

Recruitment methods

Families expecting a baby were recruited by the primary healthcare nurses during their first maternity clinic visit between 1 September 2016 and 31 December 2019. The only inclusion criterion was that the parent should speak and understand Finnish fluently. In the study area, the proportion of the Finnish-speaking population was 97% in 2018 [23]. The data collection continues until the last participating child is 1.5 years old, that is until spring 2022.

Table I. Primary aims of the prospective follow-up Central Satakunta Maternity and Child Health Clinic (KESALATU) study.

The first goal of the study is to gather comprehensive data on the use of alcohol, tobacco and other products containing nicotine in families attending maternity clinics in the Central Satakunta region.

The measurement of exhaled breath CO will be used as an objective measurement of smoking, and it will allow us to examine the effects of CO levels on maternal health during pregnancy and pregnancy outcomes.

The second goal is to examine the factors having an effect on parental smoking habits prior, during and after the pregnancy, including, for example, the role of parents' experienced stress, partner's smoking habits and development of prenatal attachment.

We will also be able to explore the importance of timing of these factors, and identify factors predisposing parents to start smoking again postpartum.

The third goal is to explore, how e.g. parental perinatal stress, prenatal smoking and intimate partner violence affect early parenthood in terms of prenatal attachment and/or capacity to mentalize the child.

The fourth goal is to explore the potential of the new pregnancy diary to support parent-foetal attachment and parental mentalization, and to decrease experienced stress and smoking behaviour.




Figure 1. Overview of the Central Satakunta Maternity and Child Health Clinic (KESALATU) Study.

Ethics

Written consent was obtained from all participants. The study data collection procedures were performed during normal maternity and child health clinic visits by the primary healthcare nurses. The participants did not receive any rewards for participating in the study. The KESALATU Study protocol was approved by The Ethics Review Committee of the Hospital District of southwest Finland. Based on the current ethical approval, it is possible to merge register data to the study data, though the permission for register data use needs to be further applied from the register-keepers.

Assessment methods

The assessment protocol included repeated (a) interview and measurement data of the mother and the child collected by maternity and child health clinic nurses and (b) parental self-report questionnaires, given to both parents (Figure 1).

Interview and measurement data. The maternity clinic nurses performed in-person interviews with the mothers and collected information about the progress of the pregnancy during the routine maternity clinic visits. During the first visit, the nurses collected

background information on maternal age, height, weight, previous pregnancies and smoking status within one year before pregnancy. After the delivery, the nurses collected information on the new-born and on pregnancy complications (pre-eclampsia, gestational diabetes, hepatogestosis and pregnancy-related blood pressure). Table II depicts the repeatedly measured and interviewed data during study.

Parental questionnaires. The questionnaires included (a) detailed questions about socio-demographic factors, (b) use of alcohol, tobacco and other nicotine products, and (c) standardized self-report questionnaires which assess the areas of specific interests: parental-foetal/infant attachment [24–27], mentalization [19,28], depression [29,30], anxiety and stress [31], and quality of life [32,33] during and after pregnancy. Table III depicts the schedule and published reliability data of the parental questionnaires and assessments.

Assessment of mentalization. Assessment of mentalization has previously only been possible using semi-structured and demanding interview methods. Recently, new self-report methods have been developed in international co-operation, which enable collecting large data [19,28]. Quality of

Table III. Published reliability data and schedule for the questionnaires in the The Central Satakunta Maternity and Child Health Clinic (KESALATU) Study assessment protocol.

Domain	Assessment method	Published reliability data	Short-term test-retest	Maternity clinic	Mid-pregnancy (27 wg)	End of pregnancy (36 wg)	After delivery check-up (6 wk postpartum)	Child health clinic	6 months postpartum	12 months postpartum	18 months postpartum	
		Coefficient alpha (internal consistency)		Early pregnancy (12 wg)				4 months postpartum				
Parental questionnaires												
Questionnaire for background information and current use of tobacco, other nicotine products, and alcohol	Heaviness of Smoking Index (HSD) [34]	0.72		M/F	M/F	M/F	M/F	M/F				
	Alcohol Use Disorders Identification Test – 3 item (AUDIT-C) [35]	0.78–0.81		F	F	F	F	M/F			M	
	Maternal/paternal antenatal attachment scale (MAAS/PAAS) [24,25]	0.82		M/F	M/F	M/F	M/F	M/F		M/F		
Mentalization	Maternal/paternal postnatal attachment scale (MPAS/PPAS) [26,27]	0.78–0.79	0.70					M/F		M/F		
	Prenatal Parental Reflective Functioning Questionnaire (P-PRFQ) [19]	0.57			M/F	M/F	M/F	M/F		M/F		
	Postnatal Parental Reflective Functioning Questionnaire (PRFQ-Fi) [28]	0.78						M/F		M/F		
Depression	Edinburgh Postnatal Depression Scale (EPDS) [29,30]		0.92	M/F	M/F	M/F	M/F	M/F		M/F		
	Perinatal Anxiety Screening Scale (PASS) [31]	0.96	0.74	M/F	M/F	M/F	M/F	M/F		M/F		
Anxiety	EUROHIS quality of life 8 item index	0.78		M/F	M/F	M/F	M/F	M/F		M/F		
	EUROHIS-QOL 8) [32]											
Routinely collected questionnaires [52]												
Everyday resources	Everyday resources for families expecting their first child				M/F							
	Everyday resources for families with a newborn baby							M/F				
	Everyday resources for families with a small child										M/F	
Violence	Domestic violence enquiry and assessment form				M/F							
	Violence enquiry for couples								M/F			

All parental questionnaires filled by both mothers and fathers. Routinely collected questionnaires filled in by mother, father, or both. F, father; mo, month; M, mother; wg weeks of gestation; wk, week.

life was measured with the EUROHIS Quality of Life 8-item index (EUROHIS-QOL 8), which is an abbreviated and translated version of the WHO-QOL-BREF [32,33]. The items are composed of overall quality of life, general health, energy, daily life activities, esteem, relationships, finances and home. Translations of the self-report measures: The Parental Reflective Functioning Questionnaire, Edinburgh Postnatal Depression Scale, EUROHIS-QOL 8, Heaviness of Smoking Index, and 3-item Alcohol Use Disorders Identification Test questionnaires have validated versions in Finnish. The Perinatal Anxiety Screen Scale and parental foetal and infant attachment questionnaires underwent an official translation/back-translation process to adapt them to the Finnish language. The back-translated versions were lastly sent to the original developers of the questionnaires, in order to contrast its conceptual content-based equivalence. The study has access also to questionnaires regarding everyday resources of the families and family violence, which are routinely collected during maternity and child health clinic visits.

Parental smoking status. Parental smoking status was first asked by the question ‘Have you smoked within one year of pregnancy?’ (no/quitted smoking/still smoking). Current smokers were asked to fill in the Heaviness of Smoking Index [34] and to report the number of cigarettes (a) per day or (b) per week. The use of e-cigarettes was asked about by the question ‘Do you use e-cigarettes daily?’. Use of alcohol during pregnancy was categorized for expectant mothers in the following way: no alcohol use/quitted when planning pregnancy/quitted when pregnancy was known/yes, current use (average doses per week). For fathers we included the 3-item Alcohol Use Disorders Identification Test [35] in all questionnaires.

The ‘Baby in Mind’ week-to-week pregnancy diary. This diary has been developed by a multidisciplinary team of clinicians and researchers from Folkhälsan, Turku University Hospital and the University of Turku, in co-operation with the Counseling Development Centre from the Department of Health and Welfare, THL. It focuses on the child’s perspective from an early stage of parenting and aims to increase the motivation to take care of one’s own and the baby’s health and well-being [17]. The ‘Baby in Mind’ pregnancy diary is available to all Finnish families expecting a baby via an electronic link [36]. A short section in the first postpartum questionnaire asks, whether and how the mother/family has used the pregnancy diary.

Exhaled breath CO

The routine use of exhaled breath CO meters began in the maternity clinics of the study area in March 2016. Thus, we have access to routine exhaled CO measurements, a biochemical marker of smoking, throughout the pregnancy. The exhaled breath CO meter offers an objective and non-invasive measurement of smoking giving instant feedback, which best suits routine maternity clinic work. Using a cut-off point of 4 parts per million (ppm) during pregnancy [37], the sensitivity of CO meters has ranged between 90 and 96%, with specificity of 92%, for those who have been smoking within 24 hours. In this study, we used the exhaled breath CO meter, the Carefusion BabyCO Monitor, which is a commercially available tool. For the study population, the CO measurement was performed three times at each visit (Table IV), and the maximum CO level was recorded. The mother was classified as non-smoker even if the CO levels were above 4 ppm if (a) the mother declared no smoking in the nurse interview despite the elevated CO levels (six women, CO level ranged from 5 to 9 ppm) and (b) self-reported no smoking in the parental assessments.

Statistical power and planned analyses of the KESALATU Study

The aim was to collect comprehensive data that covers multiple areas of interest on families during pregnancy and the first 1.5 years of offspring’s life. Information on parental smoking and the use of other nicotine products have been gathered robustly throughout the follow-up, which enables us to assess the timing of smoking and change in smoking habits. Parental health surveys were conducted repeatedly to assess changes in parental well-being. In addition, comprehensive background information on the parents was gathered in the parental questionnaires. Our rich dataset includes categorical as well as continuous variables, mainly derived from questionnaire data, which can be analysed either as a continuous variable or as a categorical variable. There are several research questions (primary aims in Table I), and multiple statistical methods will be used in the analysis of the data, thus a detailed description of each of the analyses is not possible within the scope of this paper. In the following chapters, we present the main plans briefly.

First, we will examine how the background factors of the study population differ according to smoking status, but also according to other factors, for example alcohol use stopped when planning the pregnancy v. after the positive pregnancy test result/ planned v. unplanned pregnancy/parental stress/parental mentalization capacity. We will evaluate the

Table IV. Characteristics of the Central Satakunta Maternity and Child Health Clinic (KESALATU) study population at the first trimester of pregnancy in Finnish primary healthcare in years 2016–2019.

	Pregnant women		Partners	
	<i>n</i> (%) ^a	Total <i>n</i>	<i>n</i> (%) ^a	Total <i>n</i>
Age, mean (SD)	28.9 (4.7)	247	31.4 (5.9)	156
< 25	47 (19.0)		15 (9.6)	
25–34	164 (66.4)		99 (63.5)	
35 or more	36 (14.6)		42 (26.9)	
Parity		232		
0	98 (42.2)			
1 or more	134 (57.8)			
Marital status		216		
Married/cohabiting	203 (94.0)			
Single	13 (6.0)			
Education		218		156
Under 9 years	16 (7.3)		19 (12.2)	
9–12 years	109 (50.0)		106 (67.9)	
Over 12 years	93 (42.7)		31 (19.9)	
Occupation		215		157
Full-time job	163 (75.8)		144 (91.7)	
Entrepreneur	21 (9.8)		6 (3.8)	
Part-time job	13 (6.0)		0 (0)	
Unemployed	18 (8.4)		7 (4.5)	
Yearly income of the family		194		
under 30,000 euros	30 (15.5)			
30,000–49,999 euros	51 (26.3)			
50,000–74,999 euros	81 (42.7)			
over 75,000 euros	32 (16.5)			
Smoked within one year of pregnancy	77 (31.2)	247	56 (35.4)	158
Smoking during first trimester	33 (13.4)	246	49 (31.0)	158
Number of cigarettes/day, mean (min, max)	5.7 (1, 15)	28	11.8 (1, 30)	45
Maximum CO level, mean (min, max)	6.0 (0, 23)	29		
HSI among smokers, mean (SD)	0.9 (0.9)	24	1.4 (1.3)	47
Use of e-cigarettes	2 (0.8)	245	8 (5.1)	157
Use of Swedish snuff	0 (0)	217	14 (8.9)	157
Alcohol use		217		
No alcohol use	46 (21.2)			
Quitted when planning pregnancy	51 (23.5)			
Quitted when pregnancy was known	120 (55.3)			
Yes, currently using	0 (0)			
AUDIT-C, mean (SD)			4.0 (1.9)	157
Illicit drug use	0 (0)	217	0 (0)	158

^aif not stated otherwise.

AUDIT-C, 3-item Alcohol Use Disorders Identification Test; HIS, heaviness of smoking index.

statistical differences between binary variables by using the χ^2 test. For continuous variables, independent sample *t*-test or Mann–Whitney U-test will be used depending on normality assumption.

Second, we will explore, how, for example, parental perinatal stress, smoking habits and intimate partner violence affect development of prenatal attachment and mentalization capacity. A multi-way analysis of co-variance for repeated measures will be performed to analyse the relationship between prenatal attachment and mentalization capacity and independent variables. *F* values and degrees of freedom, and 95% confidence intervals (CI) for estimated means will be

calculated. Normality assumption will be checked with a Shapiro Wilks test.

Third, factors affecting the prevalence and changes in, for example, smoking habits or the use of other nicotine containing products of the parents will be studied. For example, we will examine how the development of prenatal attachment is related to parental smoking cessation before pregnancy or at different stages of pregnancy by using multivariate logistic regression to calculate odds ratios. The relationship between parental mentalization capacity or perceived stress and changes in parental smoking habits/alcohol use can be studied similarly.

All analyses will be adjusted with appropriate confounding factors. The statistical analyses will be performed using commercially available software (SAS, version 9.4; SAS Institute Inc, Cary, NC).

The statistical power of the study was estimated by using the SAS software. Sample size calculations were formed based on the primary question regarding the association of smoking with the development of prenatal attachment and were calculated separately for mothers and fathers. We used in the calculation a *t*-test with a force of 80% and a two-sided significance level of 0.05. The primary question was if there is a difference between development of prenatal attachment for those who have smoked before/ during early pregnancy compared to non-smokers. For mothers, the standard deviation per group was estimated to be seven and for fathers to be eight, and the desired difference between groups was estimated to be five points. In this case, there should be 32 mothers per group and 42 fathers. The cohort has enough statistical power that statistically significant risk estimates can be expected for the primary questions.

Results

A total of 589 pregnant women and their partners were asked to participate in the study, and 277 (47.0%) of the pregnant women and 175 (29.6%) of the partners agreed. Twenty-nine pregnant women had an early drop-out; two due to moving outside the study area and at least 19 due to early miscarriages. Fifteen partners had an early drop-out. The final study population consisted of 248 (42.1%) pregnant women and 160 (27.1%) partners. Thirty pregnant women and two partners did not fill in the first parental questionnaire but continued in the study. Table IV depicts the characteristics of the sample during the first trimester.

The prevalence of smoking among the participating women prior to pregnancy was high (31%), but 57% of them quit smoking before the current pregnancy (Table IV). Of the partners, 35% smoked prior to the current pregnancy and only 13% succeeded in quitting smoking prior to the pregnancy. Two (1%) pregnant woman and eight (5%) partners reported using e-cigarettes during the first trimester of pregnancy. None of the pregnant women used alcohol after the pregnancy was confirmed. However, 56% had used alcohol until the pregnancy was known.

Discussion

Smoking during pregnancy is known to cause a significant risk to the course of pregnancy, maternal

health and the development of the foetus and child. It is also a risk factor that could be eliminated with more efficient prevention work. Information and awareness of this fact has long been widely available, but nonetheless, smoking during pregnancy is still too often a problem. More effective tools and intervention pathways are needed to reduce parental smoking in pregnant families and families with young children. Further, there is still sparse knowledge on both the individual differences in the psychological process of becoming a parent and the factors influencing the differences, as well as the factors influencing smoking behaviour during pregnancy, and its individual variation. This research design allows us to gain a better understanding of both these factors and their interrelationship.

The KESALATU Study is a prospective follow-up study carried out in maternity and child health clinics in primary healthcare, focusing on early parenthood and smoking behaviour. The study gathers unique research data on information: routinely collected parent and child data from the maternity and child health clinics, and study-specific data collected with repeated self-report measures and a new test battery. It covers multiple areas of interest and includes the use of some entirely new methods and tools. Together they have the potential to provide missing information on the links between early parenting psychology, risk factors, and health behaviours.

Prenatal attachment has been found to be positively associated with postnatal interaction and involvement between mother and infant [21,38], as well as child's later development [39]. There is significant individual variation in the timing and intensity of the development of prenatal attachment and mentalization. The reasons behind these individual variations are still largely unknown. It is necessary to rigorously examine the possible obstacles, such as specific psychosocial risk factors, including mental health problems or substance addictions, that make such a development of prenatal attachment and mentalization difficult or even impossible [40–42]. Early parenthood can, however, be significantly strengthened by well-targeted interventions even in the most severe risk groups [43]. One such attempt is the new pregnancy diary, which has been designed to increase the parent's curiosity towards the baby and his/her development and help the parent to think of daily situations from the baby's perspective from very early on [17].

In previous studies, the information on smoking during pregnancy has usually been based on maternal self-report, although it is known to underestimate the true prevalence of smoking [44]. The golden

standard to verify smoking status would be to use, for example, saliva or plasma cotinine [45]. In this study, maternal smoking status is verified by using the measurement of exhaled breath CO. The mean CO levels have been shown to be 5 ppm in passive and 17 ppm in active smokers [46]. It has even been suggested that exhaled CO measurement is better at measuring the consumption and changes in consumption in smoking compared to cotinine measurement [47]. This might be due to the highly variable nicotine metabolism with a half-life of 16–30 hours [48]. A recent study by Reynolds et al. [49], showed that a CO level of 3 ppm or more during the first antenatal visit associated with lower birth weight and more adverse pregnancy events. In this study, parental smoking has been assessed with repeated self-report questionnaires, with interviews conducted by nurses, and with objective measurement of the exhaled breath CO. This makes it possible to comprehensively observe and consider the effect of both the duration and amount of smoking.

Previously, it has been suggested that very few Finnish women stop smoking before pregnancy [7] because the prevalence of general smoking among women and smoking during first trimester of pregnancy have been at a similar level (Figure 1). However, our results showed that the prevalence of smoking prior to pregnancy was surprisingly high as one in three pregnant women smoked but over half of those women quit smoking prior to the current pregnancy. The use of e-cigarettes during the first trimester of pregnancy is still uncommon among pregnant women in our population, but the prevalence is likely to increase in the future as the use of e-cigarettes is increasing among young people [50]. Reports from the USA show that already 5% of women have used e-cigarettes during pregnancy [51].

A major strength of this study is that it is the first study to collect comprehensive information on early parenting psychology, prenatal psychosocial risk factors and parental health behaviour, including smoking behaviour, from standard customers of maternity and child health clinics in a primary healthcare region. The limitations of the study include rather low participation rate (42% for pregnant women and 27% for partners), which may affect the ability of the study to draw definitive conclusions. No information was collected on non-participating families. We speculate, that the longitudinal design of the study and repeated assessment of smoking by exhaled CO might have been the main reasons for declining to participate in the study. Thus, non-participating families might be in more vulnerable social situation compared to the participating families. Though our study includes comprehensive information on

parent's current education, occupation, family income, it lacks information on parents' economic stress and socioeconomic status in their childhood, which might be a precursor of smoking later on [8]. The prevalence of maternal smoking during pregnancy in this study population (13.4%) was at the same level as the smoking during early pregnancy (12–14%) in Finland (Figure 1) [6]. Thus, it seems that smoking families have participated satisfactorily in the study. In addition, the characteristics of our study population represent the standard Finnish population attending maternity clinics in terms of maternal age, parity, marital status and smoking [6], which reduces the risk of selection bias in this study.

Conducting research in primary healthcare is important in order to transmit scientific knowledge and to support the development of methods that will deliver better healthcare. Research work may stimulate the functioning of community health centres, make the personnel more committed, provide training and promote the adoption of evidence-based practices. Support from the administration of the healthcare centres and collaborative networks with specialized care and the university are important prerequisites for successful research in a primary care setting.

Acknowledgements

The authors want to thank all the nurses in the maternity and child health clinics for their valuable work recruiting the families and gathering data for this study. We also appreciate the effort of the participating families. We thank Sari Ahlqvist-Björkroth, PhD, and Prof. Anna Axelin for reviewing the Finnish versions of the attachment and perinatal anxiety questionnaires. The authors would also like to thank Prof. Susanne Somerville for permission to use the PASS questionnaire and Prof. John T. Condon for permission to use parental-foetal/infant attachment questionnaires.

Declaration of conflicting interests


The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Dr Ekblad is the chairperson of the scientific board of the Tobacco Free Finland 2030 organization. The authors declare that there is no other conflict of interests.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: this study was supported by Erkki Poikosen säätiö. Mikael Ekblad has received

the following financial support for the research: the Foundation for Paediatric Research (Lastentautien tutkimussäätiö), the Orion Research Foundation sr, the Emil Aaltonen Foundation, the Paulo Foundation, the Maud Kuistila Memorial Foundation and the Turku University Hospital Research Foundation.

ORCID iD

Mikael O. Ekblad  <https://orcid.org/0000-0001-8369-4359>

References

- [1] Knopik VS. Maternal smoking during pregnancy and child outcomes: real or spurious effect? *Dev Neuropsychol* 2009;34:1–36.
- [2] Lawder R, Whyte B, Wood R, et al. Impact of maternal smoking on early childhood health: a retrospective cohort linked dataset analysis of 697 003 children born in Scotland 1997–2009. *BMJ Open* 2019;9:e023213.
- [3] Ekblad M, Korkeila J and Lehtonen L. Smoking during pregnancy affects foetal brain development. *Acta Paediatr* 2015;104:12–18.
- [4] Ekblad M, Gissler M, Lehtonen L, et al. Prenatal smoking exposure and the risk of psychiatric morbidity into young adulthood. *Arch Gen Psychiatry* 2010;67:841–849.
- [5] Simons E, To T, Moineddin R, et al. Maternal second-hand smoke exposure in pregnancy is associated with childhood asthma development. *J Allergy Clin Immunol Pract* 2014;2:201–207.
- [6] The Finnish National Institute for Health and Welfare. Perinatal statistics – parturients, deliveries and newborns 2018. <http://urn.fi/URN:NBN:fi-fe2019121948893> (2018, accessed 22 September 2020).
- [7] Ekblad M, Gissler M, Korkeila J, et al. Trends and risk groups for smoking during pregnancy in Finland and other Nordic countries. *Eur J Public Health* 2014;24:544–551.
- [8] Lindström M, Modén B and Rosvall M. A life course perspective on economic stress and tobacco smoking: a population-based study. *Addiction* 2013;108:1305–1314.
- [9] Härkönen J, Lindberg M, Karlsson L, et al. Education is the strongest socio-economic predictor of smoking in pregnancy. *Addiction* 2018;113:1117–1126.
- [10] Villalbi JR, Salvador J, Cano-Serral G, et al. Maternal smoking, social class and outcomes of pregnancy. *Paediatr Perinat Epidemiol* 2007;21:441–447.
- [11] Finlex Data Bank. Tobacco Act of Finland, <https://www.finlex.fi/fi/laki/alkup/2016/20160549> (2016, accessed 22 September 2020).
- [12] The Finnish National Institute for Health and Welfare. Tobacco statistics, <http://urn.fi/URN:NBN:fi-fe2019121046603> (2018, accessed 22 September 2020).
- [13] Doan H and Zimmerman A. Conceptualizing prenatal attachment: toward a multidimensional view. *J Prenat Perinat Psychol Health* 2003;18:109–129.
- [14] Fava-Vizzello G, Antonioli M, Cocci V, et al. From pregnancy to motherhood: the structure of representative and narrative change. *Infant Ment Health J* 1993;13:329–336.
- [15] Pajulo M, Savonlahti E, Sourander A, et al. Prenatal maternal representations: mothers at psychosocial risk. *Infant Ment Health J* 2001;22:529–544.
- [16] Stern D. *The motherhood constellation. A unified view of parent-infant psychotherapy*. New York: Basic Books, 1995.
- [17] Pajulo H, Pajulo M, Jussila H, et al. Substance-abusing pregnant women: prenatal intervention using ultrasound consultation and mentalization to enhance the mother-child relationship and reduce substance use. *Infant Ment Health J* 2016;37:317–334.
- [18] Camoirano A. Mentalizing makes parenting work: a review about parental reflective functioning and clinical interventions to improve it. *Front Psychol* 2017;8:14.
- [19] Pajulo M, Tolvanen M, Karlsson L, et al. The prenatal parental reflective functioning questionnaire: exploring factor structure and construct validity of a new measure in the Finn brain birth cohort pilot study. *Infant Ment Health J* 2015;36:399–414.
- [20] Slade A, Patterson M and Miller M. Addendum to *Reflective functioning scoring manual* (by Fonagy et al. 1998). New York, The Psychological Center, The City College of New York, 2007.
- [21] Siddiqui A and Hägglöf B. Does maternal prenatal attachment predict postnatal mother-infant interaction? *Early Hum Dev* 2000;59:13–25.
- [22] The Finnish Ministry of Social Affairs and Health. Maternity and Child Health Clinics, <https://stm.fi/en/maternity-and-child-health-clinics> (accessed 22 September 2020).
- [23] Statistics Finland's PxWeb database, Statistics of Finland. Key figures on population by region, 1990–2018. http://pxnet2.stat.fi/PXWeb/pxweb/en/StatFin/StatFin__vrm__vaerak/statfin_vaerak_pxt_11ra.px/ (accessed 22 September 2020).
- [24] Condon JT. The assessment of antenatal emotional attachment: development of a questionnaire instrument. *Br J Med Psychol* 1993;66:167–183.
- [25] Condon JT and Corkindale C. The correlates of antenatal attachment in pregnant women. *Br J Med Psychol* 1997;70:359–372.
- [26] Condon JT and Corkindale C. The assessment of parent-to-infant attachment: development of a self-report questionnaire instrument. *J Reprod Infant Psychol* 1998;16:57–76.
- [27] Condon JT, Corkindale C and Boyce P. Assessment of postnatal paternal-infant attachment: development of a questionnaire instrument. *J Reprod Infant Psychol* 2008;26:195–210.
- [28] Pajulo M, Tolvanen M, Pyykkönen N, et al. Exploring parental mentalization in postnatal phase with a self-report questionnaire (PRFQ): factor structure, gender differences and association with sociodemographic factors. The Finn Brain Birth Cohort Study. *Psychiatry Res* 2018;262:431–439.
- [29] Cox JL, Holden JM and Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987;150:782–786.
- [30] Kernot J, Olds T, Lewis LK, et al. Test-retest reliability of the English version of the Edinburgh Postnatal Depression Scale. *Arch Womens Ment Health* 2015;18:255–257.
- [31] Somerville S, Dedman K, Hagan R, et al. The Perinatal Anxiety Screening Scale: development and preliminary validation. *Arch Womens Ment Health* 2014;17:443–454.
- [32] da Rocha NS, Power MJ, Bushnell DM, et al. The EUROHIS-QOL 8-item index: comparative psychometric properties to its parent WHOQOL-BREF. *Value Health* 2012;15:449–457.
- [33] Development of the World Health Organization WHOQOL-BREF quality of life assessment. The WHOQOL Group. *Psychol Med* 1998;28:551–558.
- [34] Borland R, Yong HH, O'Connor RJ, et al. The reliability and predictive validity of the Heaviness of Smoking Index and its two components: findings from the International Tobacco Control Four Country study. *Nicotine Tob Res* 2010;12:S45–S50.
- [35] Knibbe RA, Derickx M, Kuntsche S, et al. A comparison of the Alcohol Use Disorder Identification Test (AUDIT) in general population surveys in nine European countries. *Alcohol Alcohol Suppl* 2006;41:i19–25.
- [36] The Finnish National Institute for Health and Welfare. We're having a baby: A guidebook for expecting and looking

- after a baby, <http://urn.fi/URN:ISBN:978-952-302-833-3> (2017, accessed 22 September 2020).
- [37] Bailey BA. Using expired air carbon monoxide to determine smoking status during pregnancy: preliminary identification of an appropriately sensitive and specific cut-point. *Addict Behav* 2013;38:2547–2550.
- [38] Dubber S, Reck C, Müller M, et al. Postpartum bonding: the role of perinatal depression, anxiety and maternal–fetal bonding during pregnancy. *Arch Womens Ment Health* 2015;18:187–195.
- [39] Arguz Cildir D, Ozbek A, Topuzoglu A, et al. Association of prenatal attachment and early childhood emotional, behavioral, and developmental characteristics: a longitudinal study. *Infant Ment Health J* 2020;41:517–529.
- [40] Pajulo M, Pyykkonen N, Kalland M, et al. Substance abusing mothers in residential treatment with their babies: post-natal psychiatric symptomatology and its association with mother–child relationship and later need for child protection actions. *Nord J Psychiatry* 2011;65:65–73.
- [41] Pisoni C, Garofoli F, Tziialla C, et al. Risk and protective factors in maternal–fetal attachment development. *Early Hum Dev* 2014;90: S45–S46.
- [42] Magee SR, Bublitz MH, Orazine C, et al. The relationship between maternal–fetal attachment and cigarette smoking over pregnancy. *Matern Child Health J* 2014;18:1017–1022.
- [43] Suchman N, Pajulo M, Kalland M, et al. At-risk mothers of infants and toddlers. In: Bateman A and Fonagy P (eds) *Handbook of mentalizing in mental health practice*. Arlington: American Psychiatric Publishing, 2012, pp. 309–345.
- [44] Shipton D, Tappin DM, Vadiveloo T, et al. Reliability of self-reported smoking status by pregnant women for estimating smoking prevalence: a retrospective, cross sectional study. *BMJ* 2009;339:b4347.
- [45] Greaves R, Trotter L, Brennecke S, et al. A simple high-pressure liquid chromatography cotinine assay: validation of smoking status in pregnant women. *Ann Clin Biochem* 2001;38:333–338.
- [46] Deveci SE, Deveci F, Açık Y, et al. The measurement of exhaled carbon monoxide in healthy smokers and non-smokers. *Respir Med* 2004;98:551–556.
- [47] Secker-Walker RH, Vacek PM, Flynn BS, et al. Exhaled carbon monoxide and urinary cotinine as measures of smoking in pregnancy. *Addict Behav* 1997;22:671–684.
- [48] Benowitz NL, Hukkanen J and Jacob P III. Nicotine chemistry, metabolism, kinetics and biomarkers. *Handb Exp Pharmacol* 2009;192:29–60.
- [49] Reynolds CME, Egan B, Kennedy RA, et al. The implications of high carbon monoxide levels in early pregnancy for neonatal outcomes. *Eur J Obstet Gynecol Reprod Biol* 2019;233:6–11.
- [50] The Finnish Ministry of Social Affairs and Health. The Adolescent Health and Lifestyle Survey 2019: Adolescent tobacco and alcohol use and gambling, <http://urn.fi/URN:ISBN:978-952-00-4105-2> (2019, accessed 22 September 2020).
- [51] Kurti AN, Redner R, Lopez AA, et al. Tobacco and nicotine delivery product use in a national sample of pregnant women. *Prev Med* 2017;104:50–56.
- [52] The Finnish National Institute for Health and Welfare. Forms for counseling [Lomakkeet neuvolatyöhön] https://thl.fi/fi/web/lapset-nuoret-ja-perheet/peruspalvelut/aitiys_ ja_lastenneuvola/lomakkeet-neuvolatyohon (accessed 3 March 2021).